



Zenprox EC

KILLS: Ants, Bed Bugs, Cockroaches, Centipedes, Crickets, Flies, Fleas, Fruit Flies, Gnats, Mosquitoes, Moths, Scorpions, Silverfish, Spiders (Brown Recluse, Black Widow, others), Ticks, Wasps and Other Listed Insect Pests

FOR USE IN: Homes, Apartments, Hotels, Schools, Nursing Homes, Hospitals, Office Buildings, Warehouses, Commercial and Industrial Buildings and Other Listed Buildings and Structures

SPECIMEN LABEL

ACTIVE INGREDIENT:

Etofenprox (CAS #80844-07-1) 16.2%
Piperonyl butoxide* (CAS #51-03-6) 64.8%

OTHER INGREDIENTS*: 19.0%

TOTAL: 100.0%

* (Butylcarbityl)(6-propylpiperonyl) ether and related compounds

This formulation contains 1.4 lbs of etofenprox and 5.6 lbs of piperonyl butoxide per gallon. End use dilution is 0.25% etofenprox.

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional Precautionary Statements

EPA Reg.No. 2724-804

EPA Est. No. 2724-TX-1

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS—CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Repeated exposure to etofenprox can cause skin irritation.

FIRST AID

Call a poison control center or doctor immediately for treatment advice.

If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-248-7763 for emergency medical treatment information.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RESISTANCE-MANAGEMENT

For resistance-management, Zoecon® Zenprox® EC (Zenprox® EC) contains a Group 3A insecticide. Any insect population may contain individuals naturally resistant to Zenprox® EC and other Group 3A insecticides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly. Appropriate resistance-management strategies should be followed. To delay insecticide resistance, take the following steps:

- Rotate the use of Zenprox® EC or other Group 3A insecticides with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - o Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - o Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - o When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pests.
 - o Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - o The insect resistance-management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, record keeping and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, contact Wellmark International at 1-800-248-7763.

General Directions: For indoor use only. Zenprox® EC provides effective kill

of fleas, cockroaches and other listed pests and can be applied as an indoor broadcast, spot, void, or crack-and-crevice treatment using a low pressure sprayer, void injector or other equipment (see Specific Directions) in homes, attics and crawlspaces, apartment buildings, hotels, schools, day cares, nursing homes, hospitals, office buildings, correctional facilities, commercial and industrial buildings, warehouses, non-food/non-feed handling establishments, modes of transportation, institutions. Zenprox® EC provides up to 28 days residual control of fleas and up to 14 days residual on other listed pest species.

Apply Zenprox® EC to areas where target pests hide, such as baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, behind and under refrigerators, cabinets, sinks, furnaces, and stoves, the underside of shelves, drawers and similar areas. Pay particular attention to cracks and crevices.

Remove pets and birds and cover fish aquariums before spraying. Do not allow children or pets to contact treated surfaces until spray has dried. Do not apply to toys or stuffed animals or related articles. Do not apply this product in occupied patient rooms or in any rooms/areas while occupied by patients, the elderly, or infirm. Do not apply to classrooms while in use. Do not apply in institutions (including libraries, sports facilities, etc.) in the immediate area where occupants are present.

MIXING DIRECTIONS

Zenprox® EC mixes easily with water (diluent) and may be applied with most application equipment such as Micro-Injector or Actisol™ applicators. For best results, partially fill sprayer with water (diluent), add the labeled amount of Zenprox® EC and fill to the required amount of water (diluent). Removable chemical extraction probes (also known as “stingers”) used in suction/extraction systems must be rinsed within the pesticide container prior to removal. Diluted spray mixture may be stored overnight and used the following day, but mixture should be agitated well prior to use.

Zenprox® EC in labeled dilutions is generally non-staining and should not stain surfaces that cannot be spotted or damaged by water alone, however in questionable or sensitive surfaces, apply a small amount of dilution in an inconspicuous location to evaluate. Do not apply where electrical short circuits might result.

Do not allow to enter indoor or outdoor drains. (*No permita la entrada a desagües internos o externos.*) Follow proper disposal procedures on this label. (*Siga las indicaciones del etiquetado para el desecho apropiado del producto.*)

Do not enter or allow others to enter until spray has dried. Do not apply product in a way that will contact workers, other persons, or animals, either directly or through drift.



TANK-MIXING

Zenprox® EC can be mixed with EPA-registered insect growth regulators in areas where these products are registered to be applied. Please check product-specific label directions before using and the more stringent label directions will apply.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **Pesticide Storage:** Store in a cool, dry, and ventilated area away from humans, pets, foodstuffs, and feed. Do not store near heat or open flame. Keep from freezing. **Pesticide Disposal:** To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry). Do not pour or dispose down the drain or sewer. Call your local solid waste agency for local disposal options. **Container Handling:** Nonrefillable container. Do not reuse or refill this container. Triple rinse as follows. Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or incinerate, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SPECIFIC INSTRUCTIONS

Indoor Pests: Use Zenprox® EC as a broadcast (general surface), spot, void, or crack-and-crevice treatment to control pests inside buildings, structures, and modes of transport. This includes homes, attics and crawlspaces, apartment buildings, hotels, schools, day cares, nursing homes, hospitals, office buildings, correctional facilities, commercial and industrial buildings, warehouses, non-food/non-feed handling establishments, rail cars, marine vessels, and institutions. Zenprox® EC mixes easily with water (diluent) and may be applied with a low pressure sprayer for broadcast and spot treatments. A void injector or other application equipment that delivers low volume treatments, such as Micro-Injector or Actisol™ applicators, may be used for void and crack-and-crevice treatments.

Treatment	Specific Directions	Use Rate	Pests
Broadcast (General Surface)	Apply the dilution uniformly over surfaces to be treated. Apply to carpets, floors, cabinets, cove base, and other areas where pests are located.	0.25% Etofenprox 2 fl oz/gallon of water (diluent)/1,000 ft ²	Ants, Bed bugs, Carpet beetles, Cockroaches, Confused flour beetles, Centipedes, Crickets, Darkling beetles, Dermestids, Earwigs, Firebrats, Flies, Fleas, Fruit flies, Gnats, Indian meal moths, Lesser grain borers, Millipedes, Mosquitoes, Moths, Pillbugs, Red flour beetles, Saw-toothed grain beetles, Scorpions, Silverfish, Sowbugs, Spiders (Brown recluse, Black widow, others), Stink bugs, Ticks, Wasps
Spot, Void or Crack and Crevice	Treat entry points such as around water pipes, doors, windows, and eaves. Treat areas where pests normally feed or hide such as baseboards, corners, around water pipes, cracks and crevices. Treat void areas with sufficient material to cover the void surfaces. Spot treat floor rugs, beneath furniture, in closets, and storage areas.	0.25% Etofenprox 2 fl oz/gallon of water (diluent)/1,000 ft ²	
Spot, Void or Crack and Crevice for Bed Bugs Infesting Beds, Mattresses, Bedrooms and Sleeping Areas	Disassemble the bed frame and remove bedding. Do not apply product to bedding. Wash bedding and dry using medium to high heat to kill bed bugs. Apply to the bed frame, mattress and box spring. Application to mattresses and box springs is intended for treatment of seams, edges, and tufts only. Do not saturate the mattress and box spring. Do not treat the entire surface of the mattress or box spring. Allow spray to dry before using the mattress and/or box spring. Spot treat baseboards, behind pictures/wall hangings, rugs, and other articles in the bedroom or sleeping area.	0.25% Etofenprox 2 fl oz/gallon of water (diluent)/1,000 ft ²	Bed bugs

Repeat treatments to infested areas as necessary, but not less than 14 days after the initial treatment.

For information or in case of emergency call 1-800-248-7763 or visit our Web site: www.zoecon.com

Central Garden & Pet Company, 1501 East Woodfield Road, 200W, Schaumburg, Illinois 60173

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See product labeling for use directions.

Actisol is a trademark of Environmental Delivery Systems, Inc. Zoecon, Zoecon with design and Zenprox are registered trademarks of Wellmark International.

©2010-2026 Wellmark International. PCO 26-004

April, 2026
Schaumburg, IL